

Properties of an R plasmid pKU41 from *Pseudomonas putida* KU190

Joo Mee-Ja and Yung-Nok Lee

Department of Biology, Korea University

Pseudomonas putida KU 190에서 분리한 R plasmid pKU 41의 특성

주미자 · 이영복

고려대학교 생물학과

ABSTRACT: The properties of the plasmid pKU41 isolated from *Pseudomonas putida* KU190 have been investigated. pKU41 was defined as an R plasmid having a transmissible ampicillin and tetracycline resistance determinant, and could be classified as a plasmid belonging to IncP-1 group according to incompatibility grouping.

KEY WORDS □ R plasmid, *Pseudomonas putida*

Bacteria belonging to the genus *Pseudomonas* are able to metabolize an enormous range of natural and synthetic organic compounds, and also important agent of degradation of toxic compound that pollute the environment. The unique metabolic pathways starting from peculiar compounds in these *Pseudomonas* have required extensive genetic analysis of these bacteria. It will provide the basis for their future use as biocatalysts in chemical industries.

Studies of R factor in *Pseudomonas* has been reported in *Pseudomonas aeruginosa* as common hospital pathogen. The R factor originated from *P. aeruginosa* specified multiple drug resistance and was transferable to a variety of species (Datta *et al.*, 1971; Grinsted *et al.* 1972; Saunders and Grinsted, 1972; Olsen and Shipley, 1973). Chakrabarty *et al.* (1975) reported the transformation of *Pseudomonas putida* with an R factor DNA (RP1 plasmid). After this, a number of investigators have also reported the transformation of cell of *Pseudomonas* (Nagahari and Sakaguchi, 1978; Mercer and Loutit, 1979) and genetic analysis of

chromosomal DNA with Inc P-1 group plasmid (Hedges and Jacob, 1977; Holloway, 1978).

Gene cloning have been taken to develop cloning vectors from some of these R factors. However, most of the plasmid found in *Pseudomonas* species have high molecular weight and it makes their in vitro manipulation difficult. Plasmid pBR322 (Boliver *et al.*, 1977), one of the most widely used plasmid vector, cannot replicate in soil bacteria (Bagdasarian *et al.*, 1979) and although it was used to clone *P. aeruginosa*, some gene of *P. aeruginosa* may not be expressed in *E. coli* background (Holloway *et al.*, 1979). Therefore an appropriate cloning vector in *Pseudomonas* required. Bagdasarian *et al.* (1979) and Wood *et al.* (1981) have reported the construction of a cloning vector for *P. aeruginosa* to overcome this difficulty but they were unable to introduce it into *Pseudomonas* species. Another plasmid vector also has been developed by Ditta *et al.* (1980), which was derived from the broad-host-range plasmid RK2. Recently, host-vector systems consisting of bacterium and its vector plasmid have proven of

immense value for the cloning, amplification and analysis of DNA fragment from a wide variety of organisms. Bagdasarian *et al.* (1981) and Olsen *et al.* (1982) reported the development of host-vector system for gene cloning in *Pseudomonas*.

This paper describes some of properties of the plasmid pKU41 isolated from *P. putida* KU190. And it is one of the basic studies involved in development of a cloning vector for self-cloning of *Pseudomonas putida*.

MATERIALS AND METHODS

Bacterial strains and plasmids

The bacterial strains and plasmids used in the study are shown in table 1. *P. putida* KU190 is a strain isolated in our laboratory (Kim and Lee, 1984) and pKU41 is a cryptic plasmid isolated from *P. putida* KU190 (Kim *et al.*, 1986).

Media and growth condition

LB medium was used for all tests and growth of bacterial cells and M9 minimal medium (Maniatis *et al.*, 1982) was used for incompatibility test. *P. putida* and *E. coli* were grown with shaking at 30°C and 37°C, respectively.

Determination of antibiotic resistance.

Antibiotic resistance was determined by the procedure of Norström *et al.* (1968). The cells were grown exponentially in L-broth. The culture was diluted and spread on L-agar plates contain-

ing different concentrations of the antibiotic to be tested. Antibiotic resistance is defined as the highest concentration of an antibiotic at which 100% of the cells plated form colonies.

Curing

The method of Rheinwald *et al.* (1973) was used as follows. Cells were inoculated into 1ml of L-broth containing mitomycin C in 5 µg increment from 5 to 35 µg/ml and incubated with shaking for 36h. Then, 0.1 ml of the culture was spread on L-agar and replicated to the appropriate test plates for scoring.

Conjugation

Transfer studies was carried out by broth mating. Overnight cultures in L-broth were diluted and mixed at a donor recipient ratio of 1:10. After incubation for 60 min without shaking, the mixtures were plated directly on L-agar supplemented with the appropriate antibiotics.

Stability test

The method of sakaguchi (1981) was used. After growth in antibiotic-free medium the cells were diluted and plated on L-agar plates containing the appropriate antibiotics. Single colonies from individual plates were spotted on individual antibiotic-containing plates to score for the resistance pattern of individual colonies.

Incompatibility test

To determine the incompatibility group, The test was processed by conjugal mating with

Table 1. Bacterial strains and plasmids used.

Designation	Relevant properties (size)	References or Sources
Bacterial strains		
<i>P. putida</i> KU 190	Sal ⁺ Cat ⁺	Kim and Lee (1984)
<i>P. putida</i> TN 1032	try ⁻ leu ⁻ ben ⁻ Sm ^r rec	Nakazawa (1983)
<i>P. putida</i> TN 1307	try ⁻ leu ⁻ ben ⁻ Sm ^r rec	Nakazawa (1983)
<i>E. coli</i> HB 101	r ⁻ m ⁻ F ⁻ pro ⁻ leu ⁻ ara ⁻ gal ⁻ rec	Boyer and Roulland Dussoix (1969)
Plasmids (Inc group)		
RP4 (P-1)	Tra ⁺ Ap ^r Km ^r Te ^r (56kb)	Meyers <i>et al.</i> (1976)
RK2 (P-1)	Tra ⁺ Ap ^r Km ^r Te ^r (56kb)	Jobanputra and Datta (1974)
pMG1 (P-2)	Tra ⁺ Gm ^r Sm ^r Su ^r Hg ^r (470kb)	Hansen and Olsen (1978)
Rms 148 (P-7)	Tra ⁺ Sm ^r (145kb)	Sagai <i>et al.</i> (1976)
pKU41	(41.0kb)	Kim <i>et al.</i> (1986)

Sal⁺; salicylate utilization Cat⁺; catechol utilization Tra⁺; conjugal transfer

bacteria containing a plasmid of different Inc-group. A broth culture of KU190 was mixed with each of a series of broth cultures carrying known plasmids of different Inc-group. After incubation during the working day, each mixture were plated on L-agar supplemented appropriate drug to select for plasmid transfer.

Isolation of plasmid DNA

For detection of plasmid, the method of Kado and Liu (1981) was used. For obtaining large quantities of plasmids, the method of Tanaka and Weisblum (1975) was used. Cells were grown overnight in 25 ml of L-broth. The culture was inoculated in 1 l of L-broth and permitted to grown for 8h with shaking. The cells were harvested by low-speed centrifugation and dissolved in 10 ml of 50 mM Tris-25% (w/v) sucrose (pH 8.0), and 5 ml of lysozyme solution was added. After the mixture was kept on ice for 5 min, 4 ml of 0.25 M EDTA (Na salt, pH 8.0) was added. Also after 5 min, 5 ml of 5 M NaCl and 2 ml of 10% SDS were added and mixed. The mixture was incubated on ice for 24h, and the lysate was centrifuged. The clear supernatant was carefully decanted. To the supernatant 1/2 vol. of a polyethylene glycol solution (40% (w/v) PEG 6000) was added and mixed by inversion. The mixture was kept at 4°C overnight. The DNA precipitate was pelleted by centrifugation and redissolved in TE buffer (0.05 M Tris-0.02 M EDTA, pH 8.0). Then 8.0 g of CsCl and 40 ml of an etidium bromide solution (1% (w.v)) were added. The density of the solution was adjusted to 1.60 g/ml by addition of a small amount TE buffer. Ultracentrifugation was carried out in an RP 65T rotor at 38,000 rpm and for 48h, Under ultraviolet illumination, plasmid DNA was collected, and was extracted three times with isopropyl solution saturated NaCl. The plasmid DNA solution was dialized against three changes of TE buffer.

Transformation

The procedure described by Nagkazawa (1983) was used for transformation. Cells were grown overnight in L-broth with shaking. A portion (1.5 ml) of such a culture was inoculated in 20 ml of 0.5 M sucrose in L-broth and grown for 3-4h. The cells were harvested by centrifugation and washed

with 10 ml of 10 mM NaCl in 0.5 M sucrose solution. The cells were again centrifuged, and resuspended in 10 ml of 50 mM CaCl₂ in 0.5 M sucrose solution. The cells were kept ice for 20 min, harvested by centrifugation and resuspended in 2 ml of 50 mM CaCl₂. 0.2 ml of these cells was then mixed with 0.1 ml of DNA solution and incubated at 0°C for 60 min. The cell DNA mixture was then subjected to a heat pulse at 42°C for 2 min, chilled, and finally diluted in 10 volume of fresh L-broth. The cells were allowed to grown at 30°C on a shaker and aliquots were plated on L-agar plates containing the appropriate antibiotics.

Electrophoresis

Electrophoresis of DNA was carried out in agarose gel as described by Casse *et al.* (1979). Sample were run in 0.8% agarose slab gels in a horizontal electrophoresis apparatus filled with electrophoresis buffer (40 mM Tris-acetate, 20 mM sodium acetate and 1 mM Na₂-EDTA). Samples of DNA were mixed with 0.015% bromophenol blue in 60% sucrose, 60 mM EDTA. Electrophoresis was performed at 120 V for 3-4h.

RESULTS AND DISCUSSION

Antibiotic resistance of *P. putida* KU190

Antibiotics used were ampicillin, carbenicillin, chloramphenicol, gentamycin, kanamycin, rifampicin and streptomycin. *P. putida* KU190 was resistant to ampicillin, carbenicillin, chloramphenicol and tetracyclin whereas sensitive to gentamycin, kanamycin, rifampicin and streptomycin (Table 2).

Curing of pKU41

Curing is one of the main criteria for determining character encoded by plasmid. The strain treated with mitomycin C (25 ug/ml) lost resistant to ampicillin, carbenicillin and tetracyclin (Table 3). Accordingly, resistance to these antibiotics is specified by pKU41, and pKU41 was defined as an R plasmid having the resistance-determining gene.

Transfer ability of pKU41

Conjugal mating was carried out in order to

Table 2. Resistance of *P. putida* KU 190 to various antibiotics

Antibiotics	Concentration of antibiotics ($\mu\text{g}/\text{ml}$)						
	25	50	100	200	400	800	1600
Ap	+	+	+	+	+	+	-
Cb	+	+	+	+	+	+	+
Cm	+	+	+	+	+	+	+
Gm	-	-	-	-	-	-	-
Km	-	-	-	-	-	-	-
Rif	+	-	-	-	-	-	-
Sm	+	-	-	-	-	-	-
Tc	+	+	-	-	-	-	-

Ap, ampicillin; Cb, carbenicillin; Cm, chloramphenicol; Gm, gentamycin; Km, kanamycin; Rif, rifampicin; Sm, streptomycin; Tc, tetracycline; +, growth; -, no growth

Table 3. Effect of mitomycin C (25 $\mu\text{g}/\text{ml}$) on curing

Selective marker	Concentration of antibiotics ($\mu\text{g}/\text{ml}$)	Frequency of curing (%) ^a
Ap	800	21.4
Cb	1600	29.4
Tc	50	16.0


a; No. of cured colonies/total No. of examined colonies

study the transfer ability of the plasmid pKU41. The transconjugants was obtained from *P. putida* TN1307, TN1032 and *E. coli* HB101 at the frequency between 10^{-6} - 10^{-5} (Table 4). Therefore pKU41 can be transferred to *E. coli* as well as

Table 4. Transfer frequency of pKU41 by mating.

Donor	Recipient	Selective marker (conc. of marker)	Transfer frequency ^a
KU 190 (pKU41)	<i>P. putida</i> TN 1307	Ap (800 $\mu\text{g}/\text{ml}$)	2.5×10^{-5}
		Tc (25 $\mu\text{g}/\text{ml}$)	
		Sm (100 $\mu\text{g}/\text{ml}$)	
	<i>P. putida</i> TN 1032	Ap (800 $\mu\text{g}/\text{ml}$)	1.1×10^{-5}
		Tc (25 $\mu\text{g}/\text{ml}$)	
		Sm (100 $\mu\text{g}/\text{ml}$)	
	<i>E. coli</i> HB 101	Ap (800 $\mu\text{g}/\text{ml}$)	1.0×10^{-6}
		Tc (25 $\mu\text{g}/\text{ml}$)	
		Sm (100 $\mu\text{g}/\text{ml}$)	

a: No. of transconjugants/No. of donor cells

**Fig. 1.** Agarose gel electrophoresis of transconjugants.

lane 1; wild type, KU190 (pKU41)
lane 2; *P. putida* TN1307 (pKU41)
lane 3; *P. putida* TN1032 (pKU41)
lane 4; *E. coli* HB101 (pKU41)

strains of *P. putida* (Fig. 1). And it is one of properties of IncP-1 group plasmid.

Stability of pKU41

Studies of stability of pKU41 was carried out using antibiotic resistance of transconjugant acquired the plasmid. pKU41 was very stable in both *P. putida* TN1307 (98.0%) and *E. coli* HB101 (97.5%) as in the case of wild strain (99.7%) (Table 5).

Table 5. Stability of pKU41 in different strains

Host strains	Selective marker ^a	Stability(%) ^b
<i>P. putida</i> KU190 (pKU 41)	Ap, Tc	99.7
<i>P. putida</i> TN1307 (pKU 41)	Ap, Tc	98.0
<i>E. coli</i> HB101 (pKU 41)	Ap, Tc	97.5

^a; The concentration of ampicillin (Ap) and tetracycline (Tc) were 600 $\mu\text{g}/\text{ml}$ and 20 $\mu\text{g}/\text{ml}$

^b; Stability is indicated as the percentage of cells resistant to ampicillin and tetracycline after one generation in antibiotic free L-broth

Table 6. Incompatibility of pKU41 for different plasmids.

Donor	Recipient	Plasmid (Inc. group)	Selective marker	Conc. of marker	No. of colonies on master plate
KU190 (pKU41)	<i>E. coli</i>	RP ₄ -RK ₂	Km	50 µg/ml	N.D
	C 600	(p-1)	Ap Tc	800 µg/ml 25 µg/ml	
	<i>P. aeruginosa</i>	pMG1	Gm	25 µg/ml	5.2X10 ²
	PAO 303	(p-2)	Ap	800 µg/ml	
			Tc	25 µg/ml	
	<i>P. aeruginosa</i>	Rms 148	Sm	100 µg/ml	5.2X10
PAO 303	(p-7)	Ap	800 µg/ml		
		Tc	25 µg/ml		

ND; Not detected.

Incompatibility of pKU41

Incompatibility is currently the basis of plasmid classification. Therefore, nearly all plasmids may be assigned to one of a relatively small number of such groups. The incompatibility properties of pKU41 were tested by conjugal mating and the results was scored as the colonies retaining determinants carried by the income and resident plasmid; A high percent figure therefore indicates compatibility, a low figure incompatibility. pKU41 was seem to be compatible with pMG1 as P-2 and Rms 148 as P-7 and incompatible with RP₄ and RK₂ as P-1 (Table 6). This suggests that pKU41 fall therefore into the same incompatibility group as RP₄ and RK₂.

Transformation of *P. putida* and *E. coli* with pKU41

Transformation exhibit a spectrum of host expressing gene of plasmid. Although the frequency of transformants obtained differs for *P. putida*

(2.8x10⁻⁵) and *E. coli* (7.8x10⁻⁷), transformant containing pKU41 could be obtained from the two strains (Table 7). And the presence of pKU41 in




Fig. 2. Agarose gel electrophoresis of transformants. lane 1; wild type. KU190 (pKU41) transformants
lane 2; *P. putida* TN1307 (pKU41)
lane 3; *E. coli* HB101 (pKU41)

Table 7. Transformation frequency for antibiotic resistance characters specified by pKU 41.

Donor plasmid	Recipient	Antibiotic maker	Selective conc.	Transformation frequency
pKU41	<i>P. putida</i>	Ap	800 µg/ml	2.8x10 ⁻⁵
		TN 1307	Tc 25 µg/ml	
	<i>E. coli</i>	Ap	800 µg/ml	7.8x10 ⁻⁷
		HB 101	Tc 25 µg/ml	

Transformation frequency: transformant No./µg DNA/viable cells

both strain led to cellular resistance to ampicillin and tetracyclin (Fig. 2). It suggests that pKU41 can be used to transform both *P. putida* and *E. coli* strain to antibiotic-resistance characters.

As stated in the introduction, R plasmid originated *P. aeruginosa* such RP1 and RP4 are well documented, there is so far no report concerning an R plasmid isolated from *P. putida* as soil saprophytes. However, Grinsted *et al.* (1972) indicated the base composition of an R factor obtained from *P. aeruginosa* (RP1) is more characteristic

of terrestrial and aquatic saprophytic *Pseudomonas* than *P. aeruginosa*. Thus terrestrial saprophytic *Pseudomonas* may be the natural hosts for R factor.

The occurrence of pKU41 as R plasmid in *P. putida* may be extremely useful because pKU41 having a transmissible and ampicillin and tetracycline resistance determinant would be of value in an attempt to development for a cloning vector in *Pseudomonas*.

적 요

Pseudomonas putida KU 190에서 분리한 plasmid pKU 41의 특성을 조사하였다. pKU 41은 전달성 및 암피실린과 테트라사이클린에 내성을 나타내는 유전자를 지닌 R. plasmid인 것으로 밝혀졌으며, 불화합성에 따르는 분류로는 IncP-1군인 것으로 나타났다.

REFERENCES

1. Bagdasarian, M., M.M. Bag dasarian, S. Coleman, and K. Timmis, 1979. New vector plasmids for gene cloning in *Pseudomonas*. *Dev. Genet.* **1**, 411-422.
2. Bagdasarian, M., R. Lurz, B. Brückert, F.C.H. Franklin, M.M. Bagdasarian, J. Forrey and K.N. Timmis, 1981. Specific-purpose plasmid cloning vectors II. Broad host range, high copy number. RSF 1010-derived vectors and a host-vector system for gene cloning in *Pseudomonas*. *Gene* **1**, 237-247.
3. Boliver, R., R. Rodriguez, P. Greene, M. Betlach, H. Heyneker, H. Boyer, J. Crosa and S. Falkow, 1977. Construction and characterization of new cloning vehicles, II. A multipurpose cloning system. *Gene (Amst.)* **2**, 95-113.
4. Boyer, H.W., and D. Roulland-Dussoix, 1969. A complementation analysis of the restriction and modification of DNA in *Escherichia coli*. *J. Mol. Biol.* **41**, 459-472.
5. Casse, F., C. Boucher, J.S. Julliot, M. Michel, and J. Denarie, 1979. Identification and characterization of large plasmids in *Rhizobium meliloti* using agarose gel electrophoresis. *J. Gen. Micro.* **113**, 229-242.
6. Chakrabarty, A.M., 1975. Transformation of *Pseudomonas putida* and *E. coli* with plasmid-linked drug-resistance factor DNA. *Proc. Natl. Acad. Sci. USA.* **72**, 3647-3651.
7. Datta, N., R.W. Hedpes, E.J. Shaw, R.B. Sykes and M.H. Richmond, 1971. Properties of an R factor from *Pseudomonas aeruginosa*. *J. Bacteriol.* **108**, 1244-1249.
8. Ditta, G., S. Stanfield, D. Corbin, and D. Helinski, 1980. Broad host range DNA cloning system for gram-negative bacteria; construction of a gene bank of *Rhizobium meliloti*. *Proc. Natl. Acad. Sci. USA.*, **77**, 7347-7351.
9. Grinsted, J., J.R. Saunders, L.C. Ingram, R.B. Sykes and M.H. Richmond, 1972. Properties of an R factor which originated in *Pseudomonas aeruginosa* 1822. *J. Bacteriol.* **110**, 529-537.
10. Hansen, J.B. and R.H. Olsen, 1978. Isolation of large plasmid and characterization of P2 incompatibility group plasmids pMG1 and pMG5. *J. Bacteriol.* **135**, 227-238.
11. Hedges, R.W., and A.E. Jacob. 1977. Wide ranging plasmid bearing the *Pseudomonas aeruginosa* tryptophan synthetase genes.

- Nature. 267, 283-284.
12. Holloway, B.W. 1978. Isolation and characterization of an R'plasmid in *Pseudomonas aeruginosa*. *J. Bacteriol.* 133, 1078-1082.
 13. Holloway, B.W., V. Krishnapillai, and A.F. Morgan. 1979. Chromosomal genetics of *Pseudomonas*. *Microbiol. Rev.* 43, 73-102.
 14. Jobanputra, R.S. and N. Datta, 1974. Trimethoprim R factors in *Enterobacteria* from clinical specimens. *J. Med. Microbiol.* 7, 169-177.
 15. Kado, C.I. and S.T. Liu, 1981. Rapid procedure for deletion and isolation of large and small plasmid. 145, 1365-1373.
 16. Kim, H.K., S.K. Koh and Y.N. Lee, 1986. Restriction map of a cryptic plasmid from *Pseudomonas putida*, *Kor. J. Microbiol.* 24, 7-11.
 17. Kim, J.K. and Y.N. Lee, 1984. Isolation and identification of *Pseudomonas* utilizing hydrocarbon. *Kor. J. Microbiol.* 22, 29-34.
 18. Maniatis, T., E.F. Fritsch and J. Sambrook (eds.), 1982. *Molecular cloning; A Laboratory manual*, Cold Spring Harbor Laboratory. New York.
 19. Mercer, A.A. and J.S. Loutit, 1979. Transformation and transfection of *Pseudomonas aeruginosa*: effects of metal ions. *J. Bacteriol.* 140, 37-42.
 20. Meyers, J.A., D. Sanchez, L.P. Elwell, and S. Falkow, 1976. Simple agarose gel electrophoresis method for the identification and characterization of plasmid deoxyribonucleic acid. *J. Bacteriol.* 127, 1529-1537.
 21. Nakazawa, J. 1983. 遺傳子組換え實用化 技術 第四集 V. シコードモナス, 73-85.
 22. Nordström, K., K.G. Eriksson-Greenberg and H.G. Boman, 1968 Resistance of *Escherichia coli* to penicillins, III. AmpB, a locus affecting episomally and chromosomally mediated resistance to ampicillin and chloramphenicol. *Genet. Res.* 12, 157-168.
 23. Olsen, R.H. and P. Shipley, 1973. Host range and Properties of the *Pseudomonas aeruginosa* R factor R1822. *J. Bacteriol.* 113, 772-780.
 24. Olsen, R.H., G. DeBusscher, and W.R. McCombie, 1982. Development of broad-host-range vectors and gene banks: self-cloning of the *Pseudomonas aeruginosa* PAO chromosome. *J. Bacteriol.* 150, 60-69.
 25. Rheinwald, J.G., A.M. Chakrabarty, and I.C. Gunsalus, 1973. A transmissible plasmid controlling camphor oxidation in *Pseudomonas putida*. *Proc. Nat. Acad. Sci. USA* 70, 885-889.
 26. Sagai, H., K. Hasuda, S. Iyobe, L.E. Bryan, B.W. Holloway and S. Mitsuhashi, 1976. Classification of R plasmids by incompatibility in *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* 10, 573-578.
 27. Sakaguchi, K., 1981. Vectors for gene cloning in *Pseudomonas* and their applications. *Current topics in microbiology and immunology.* 96, 31-45.
 28. Saunders, J.R. and J. Grinstead. 1972. Properties of RP4, an R factor which originated in *Pseudomonas aeruginosa* S8. *J. Bacteriol.* 112, 690-696.
 29. Tanaka, T. and B. Weisblum, 1975. Construction of a colicin EI-R factor composite plasmid in vitro: Means for amplification of deoxyribonucleic acid. *J. Bacteriol.* 121, 354-362.
 30. Wood, D.O., M.F. Hollinger, and M.B. Tindol, 1981. Versatile cloning vector for *Pseudomonas aeruginosa*. *J. Bacteriol.* 14, 1448-1451.

(Received Apr. 28, 1987)