

MINIREVIEW

Biodegradation of Dibenzo-*p*-dioxin and Dibenzofuran by Bacteria

Jean Armengaud* and Kenneth N. Timmis

Department of Environmental Microbiology, GBF-National Research Centre for Biotechnology, Mascheroder Weg 1, 38124 Braunschweig, Germany

Polychlorodibenzofurans and polychlorodibenzo-*p*-dioxins are among the most toxic xenobiotics released into the biosphere and the cause of significant public concern because of their apparent ubiquity - albeit at low levels - in food and environment. Several bacteria able to degrade nonchlorinated dioxin and dibenzofuran and in some cases to attack chlorinated analogues have recently been isolated. This opens up the possibility that bioremediation processes may ultimately be developed to eliminate these toxic compounds from contaminated sites. In this review we summarize current knowledge on the genetics and biochemistry of dioxin and dibenzofuran degradation by *Sphingomonas* sp. RW1, a gram-negative bacterium, and highlight the unusual nature of the genetic organization of these pathways.

Biodegradation of dibenzofuran and dibenzo-*p*-dioxin

Aromatic and aliphatic hydrocarbons, and particularly synthetic derivatives thereof, constitute major environmental pollutants. Microorganisms are major agents of carbon mineralization and are largely responsible for eliminating such compounds from the biosphere. They are, however, unable or poorly able to catabolize effectively the more toxic industrial pollutants that have entered the environment. The effective exploitation of microbial catabolic activities in biotechnological processes to detoxify and destroy the more recalcitrant environmental pollutants requires that we build a knowledge base on relevant catabolic processes in the environment, and the ecological factors which influence their efficiency, so that rational strategies to optimize required activities can be developed

from an understanding of the critical metabolic processes.

Microbial degradation of monocyclic aromatics such as benzene, toluene and their chloroderivatives (Table 1) has been intensively studied (73), and the various metabolic strategies and mechanisms exploited by microorganisms in the utilization of these compounds as carbon and energy sources have been the subject of numerous reviews (30, 55). More recently, microbial metabolism of bicyclic and polycyclic aromatics, such as polychlorinated biphenyls (PCB), dibenzo-*p*-dioxins (DBD), dibenzofurans (DBF) and polyaromatic hydrocarbons (PAH), have received increasing attention (1, 67). Whereas microbial transformations of PCBs and PAHs have been characterized extensively (5, 24, 27, 32, 33, 37, 38), and research currently focusses on improvement of the catabolic properties of the enzymes and direct applications (24, 41, 63), progress on microbial transformation of DBF and DBD has been slow due to the difficulty of isolating microorganisms able to attack these compounds, the lack of commercially available, environmentally-relevant chlorinated congeners, and the problems of studying these highly toxic compounds. It is well known that some DBD congeners, such as 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD), are extremely toxic: TCDD is 500 times higher than cyanide and the ingestion of 1 mg/kg is lethal for adult humans. TCDD is also cancerogenic and teratogenic. Its high chemical stability and hydrophobicity make it very persistent and difficult to remove from contaminated sites (25).

Chlorinated DBDs and DBFs are created as byproducts or trace contaminants in a variety of chemical processes (manufacture of herbicides, insecticides and fungicides, paper pulp bleaching and metal smelting), as well as in combustion processes (burning of solid wastes in incinerators, motor vehicle exhausts and accidental fires). The 1976 Séveso example in which an accident in a chemical plant manufacturing the herbicide 2,4,5-trichlorophenol

* To whom correspondence should be addressed. Phone: 49-531-6181-403. Fax: 49-531-6181-411. E-mail: jar@gbf-braunschweig.de.




Fig. 1. Proposed catabolic pathways for dibenzofuran and dibenzo-*p*-dioxin in *Sphingomonas* sp. RW1. Chemical designations are (I) dibenzo-*p*-dioxin, (II) dibenzo-*p*-dioxin-*cis*-dihydrodiol, (III) 2,2',3-trihydroxydiphenyl ether, (IV) 2-hydroxy-6-oxo-6-(2-hydroxyphenyl)-hexa-2,4-dienoate, (V) catechol, (C6) 2-hydroxy-*cis*,*cis*-muconate, (VI) dibenzofuran, (VII) dibenzofuran-*cis*-dihydrodiol, (VIII) 2,2',3-trihydroxybiphenyl, (IX) 2-hydroxy-6-oxo-6-(2-hydroxyphenyl)-hexa-2,4-dienoate, (X) salicylate, (C5) 2-hydroxypentadienoate, (XI) 2-hydroxymuconate semialdehyde, (XII) 2-hydroxypenta-2,4-dienoate, (XIII) 4-hydroxy-2-oxovalerate and (XIV) gentisate. Unstable compounds which spontaneously transform to other products are indicated in brackets.

which is then transformed by an aldolase to products that enter the Krebs cycle (Fig. 1). Such a pathway is expected to occur in *Sphingomonas* sp. RW1.

Some of the enzyme of these pathways have now been purified and characterized biochemically, and in some cases genetically. First, we shall present

The initial dioxygenase is a three-component enzyme with an atypical electron supply system

The stereospecific angular dioxygenation of dibenzofuran and dibenzo-*p*-dioxin is carried out by a dioxin dioxygenase, which has been purified from RW1 and characterized by Bünz and Cook (13). Crude cell extracts exhibited relatively low levels of enzyme which was unstable (80% loss of activity after storage at 4°C for 24 h). A nonlinear activity: protein concentration relationship was observed, suggesting that the enzyme is a multicomponent complex. The properties of the five polypeptides purified by Bünz and Cook (13) are presented in Table 1. In fact, the enzymatic complex consists only of four of these polypeptides, as is the case for other three-component dioxygenases acting as ring hydroxylases of monocyclic or bicyclic aromatics (15). The initial dioxygenase itself is a $\alpha_2\beta_2$ heterodimer of two of these polypeptides, which requires electrons in order to ring hydroxylate its substrate. These electrons derive from NADH plus H⁺ which is oxidized to NAD⁺ by a specific reductase, which in turn transfers them to a ferredoxin, which then reduces the large subunit of the initial dioxygenase (Fig. 2).

Two reductases (components A1 and A2), which are monomeric flavoproteins of about 44 kDa containing a labile flavin adenine dinucleotide cofactor, were purified and shown to be able to reduce the ferredoxin. Neither reductase contained an [Fe-S] cluster which clearly distinguishes them from reductases associated with Class III ring hydroxylating oxygenases, typified by the naphthalene dioxygenase (22) and from those associated with Class IA and Class IB enzymes (9). The N-terminal sequences of these two proteins were determined by Edman degradation (13); no significant similarity to other known flavoproteins was detected using these sequences as queries. The electron transport system associated with the dioxin dioxygenase therefore appears unusual.

The electron transfer activities of the reductases could be easily measured by reduction of some redox indicators such as dichlorophenolindophenol, or in a coupled reaction with the ferredoxin by a cytochrome C reduction assay. The two reductases were found to be isofunctional as they could both reduce the ferredoxin which transfers electrons to the dioxin dioxygenase in an *in vitro* reconstituted




Fig. 2. The dioxin dioxygenase and its electron supply system. The reaction carried out by the multicomponent ring-hydroxylating dioxin dioxygenase and its electron transfer chain is shown. A flavoprotein reductase, RedA2, accepts electrons from NADH and transfers them *via* the ferredoxin Fdx1 to the terminal oxygenase. The reduced terminal oxygenase catalyses the angular oxidation of dibenzo-*p*-dioxin and some other aromatic substrates.

the A1 reductase in RW1 cells grown under inducing conditions for the dioxin dioxygenase. These aspects favour the A2 flavoprotein as the main (but not necessarily exclusive) reductase constituent of the electron transfer chain of the dioxin dioxygenase.

The second component of the redox chain is a 12 kDa protein which exhibits all the usual characteristics of ferredoxins: small size, acidic behaviour and presence of an [Fe-S] prosthetic group (12). From the N-terminal sequence determination of the purified ferredoxin, it was deduced that it shares some similarities with a group of electron carriers comprising the putidaredoxin of *Pseudomonas putida* (42, 54), a ferredoxin from a photosynthetic bacterium (3, 50), a ferredoxin from *Escherichia coli* (65) and the mammalian adrenodoxin (16). All these ferredoxins contain a [2Fe-2S] cluster coordinated in the polypeptide by four cysteine residues arranged in the pattern Cys-Xaa₅-Cys-Xaa₂-Cys-Xaa_n-Cys. Some of these proteins have been identified as the electron donors for monooxygenases involved in the catabolism of aromatic compounds, namely FdxP from *Caulobacter crescentus* (69), the terpredoxin from *Pseudomonas* sp. (53), the putidaredoxin from *Pseudomonas putida* (54) and the rhodocoxin from *Rhodococcus* sp. strain NI86/21 (48), which are involved in the hydroxylation of *p*-hydroxybenzoate, α -terpineol, camphor and S-ethyldipropylcarbamothioate, respectively. From a phylogenetic point of view, these ferredoxins are unrelated to the Fe-S proteins which usually function as electron donors to Class IIB ring hydroxylating dioxygenases and which contain a Rieske-type [2Fe-2S] cluster coordinated by two cysteines and two histidines. It has therefore been

proposed (13) that the new multiple component dioxin dioxygenase of *Sphingomonas* sp. RW1 is a class IIA ring hydroxylating dioxygenase, according to the classification scheme of Batié *et al.* (9) which is essentially based on the characteristics of the electron supply system. The dioxin dioxygenase is the second representative of this class, the first one being the pyrazon dioxygenase described by Sauber *et al.* (56).

In the case of the dioxygenase itself, the size of the two components, 45 kDa for the large subunit and 23 kDa for the small subunit, as well as the presence of a Rieske-type [2Fe-2S] cluster in the large subunit as revealed by ultra-violet/visible spectroscopy of the purified native protein, indicate that the structure of the dioxin dioxygenase is similar to that observed for other three-component dioxygenases (15). However, the 20 amino acid N-terminal sequence of the α subunit of the dioxin dioxygenase does not exhibit any significant similarities with other known dioxygenases (Table 1).

The activity of the dioxin dioxygenase was assayed in a reconstituted *in vitro* system. It transformed dibenzofuran to trihydroxybiphenyl, and dibenzo-*p*-dioxin to trihydroxydiphenyl ether, with concomitant consumption of one molecule of oxygen per molecule of product formed. Therefore, it appears that the same enzymatic complex initiates the catabolic pathways of both substrates (Fig. 1) in RW1. In contrast to all other known multi-ring dioxygenases, such as the toluene dioxygenase (30) and the biphenyl dioxygenase (5, 23), *in vitro* the purified dioxin dioxygenase specifically attacks the dibenzo-*p*-dioxin molecule by introducing two atoms

of oxygen at a pair of vicinal carbon atoms, one of which is involved in one of the bridges between the two aromatic rings, rather than the neighboring carbon atoms not involved in the bridges (13). These results confirmed the first descriptions of this angular attack presented by Engesser *et al.* (21) and by Wittich *et al.* (71). In addition, the dioxin dioxygenase can attack biphenyl, xanthene, fluoren-9-one and dibenzothiophene. In the latter case, the products of the reaction are presumably the sulfoxide and the sulfone of dibenzothiophene. As one of the major sources of acid rain is sulphur emission from coal combustion (31), the microbial degradation of this organosulphur compound is of interest for the potential desulphurization of coal and fossil fuels.

The dioxin dioxygenase complex appears to be very unstable due to its sensitivity to oxygen and temperature. Its purification and characterization by Bünz and Cook (13) therefore represents a significant advance and has provided important insights into its structure and activity.

Characterization of the electron transport system associated with the dioxin dioxygenase

Initial efforts to genetically analyse the dioxin catabolic pathway of RW1 met with only limited success which was restricted to the readily scored *meta*-cleavage enzymes. However, the purification of the polypeptides of the dioxin dioxygenase, the determination of their N-terminus sequences and the growing database of sequences of degradative enzymes, opened the way for new strategies. Sequence similarity searches and alignment of related proteins reveal consensus sequences in functionally critical regions. This in turn permits the design of PCR amplification strategies to generate specific oligonucleotide probes for the corresponding structural genes, for example with a degenerate forward primer based on the experimentally-determined N-terminal sequence of the purified protein and a degenerate reverse primer based on a consensus sequence found in its C-terminal region (Fig. 3A). Such a strategy was used to clone the genes specifying the reductase A2 and the ferredoxin. However, in the latter case, no well-conserved sequence at the C-terminus of this class of small proteins is found. Comparison of the sequences of the four most closely-related ferredoxins revealed, however, a stretch of six amino acids, four of which are conserved in all the proteins. From this sequence, a primer was designed which was short (17 nt) and highly degenerate (4). The PCR strategy was therefore improved by addition of a second, more

selective, amplification step using a third primer.

This nested-PCR strategy is schematically presented in Fig. 3B. It involved the generation of a rather unspecific group of PCR fragments using one forward primer based on the N-terminal sequence of the protein and the highly degenerate primer based on the short conserved C-terminal sequence, the isolation of these PCR fragments, and the use of these as a template for a second PCR amplification with the same reverse primer and a new forward primer corresponding to a second re-




Fig. 3. PCR-based strategy developed to clone the genes involved in the dibenzofuran and dibenzo-*p*-dioxin pathways. Panel A: The N-terminal sequence of each purified polypeptide (a), determined by Edman degradation, was aligned with the sequences of similar polypeptides (b). From this alignment, a C-terminal consensus sequence was defined. A degenerate forward primer was then designed from the N-terminal consensus sequence (AJ01) and a degenerate reverse primer from the consensus C-terminal sequence (AJ02). Panel B: To render more specific the amplification of the desired fragment, a third primer can be used in a nested-PCR reaction. Degenerate PCR primers AJ01 and AJ03 were designed according to the N-terminal sequence of the protein while the reverse-primer AJ02 is designed from a C-terminal consensus sequence. The first PCR reaction is carried out with the AJ01 and AJ02 primers using total genomic DNA from the bacterial strain, giving a mixture of DNA fragments as a result of the high degeneracy of the primers. After purification, these products are used as template in a second, more specific reaction using degenerate primers AJ03 and AJ02.

gion of the known N-terminal sequence of the protein. This yielded a specific probe for the ferredoxin, designated *fdx1*, which was then used to screen a previously constructed pLAFR3-based cosmid library of RW1 DNA (34). A 4.6 kb DNA fragment encompassing the ferredoxin gene was thereby identified (4). Whereas the genes encoding the α and β subunits of multi-component dioxygenases have thus far always been found to be contiguous with the gene of the cognate electron carrier (Fig. 4), the *fdx1* gene of RW1 is not linked with the dioxin dioxygenase genes. Rather, it is clustered with genes apparently encoding two atypical decarboxylases/isomerases and a glutathione S-transferase (Fig. 5). A role for such proteins in the proposed di-

benzo-*p*-dioxin and dibenzofuran degradative pathways of RW1 (71) has not thus far been suggested, so their functions and possible involvement in these pathways is presently unclear. It may be noted, however, that glutathione S-transferase genes are often found in operons encoding catabolic pathways for multiring aromatic compounds, as exemplified by the *bphK* gene of *Burkholderia* LB400, a strain able to degrade biphenyl and some polychlorinated biphenyls (Hofer *et al.*, 1994), although the functional role of such enzymes in these pathways remains to be established (68).

In order to verify that the *fdx1* gene encodes a functional electron carrier, it was hyperexpressed in *Escherichia coli* and the resulting protein was pu-




Fig. 4. Usual genetic organization of three-component dioxygenases. The organisation of the genes encoding the different polypeptides of representative hydroxylating enzymes is shown. The general name of each cluster, the name of the microorganism containing the enzymes, as well as the class (according to Batie *et al.* [9]) to which they belong, are indicated in front of the clusters. Gene names are given above each ORF and the arrows indicate the direction of transcription. Each cluster was aligned with the others, taking the gene specifying the α subunit dioxygenase as a reference. Genes are drawn according to their size and relative position. Abbreviations for genes and references are: *cbdABC* for plasmid-encoded 3-chlorobenzoate-4,5-dioxygenase of *Alcaligenes* sp. BR60 (49), *xylXYZ* for benzoate and toluate dioxygenase of *Pseudomonas putida* mt-2 (51), *todC1C2BA* for toluene dioxygenase of *P. putida* F1 (72), *bphA1A2A3A4* for biphenyl dioxygenase from *Rhodococcus globerulus* P6 (5), *bphGAEF* for biphenyl dioxygenase from *Comamonas testosteroni* sp. B356 (64), *bphAEFG* for biphenyl dioxygenase from *Burkholderia* sp. LB400 (23), *doxABD* for dibenzothiophene dioxygenase from *Pseudomonas* sp. (17), *nahAabcd* for naphthalene dioxygenase from *Pseudomonas* sp. strain 9816-4 (Parales, unpublished), and *ntdAabcd* for 2-nitrotoluene dioxygenase from *Pseudomonas* sp. JS42 (52).




Fig. 5. Genetic organization of the locus encompassing the ferredoxin gene. The position and orientation of the different ORFs detected within the 4.5 kb *Bam*HI fragment are shown by full arrows. Transcription of the genes is from left to right. A scale bar on the bottom of the figure indicates the absolute gene sizes in kilobase pairs (kb). ORF1 and ORF2 encode two putative decarboxylases whereas ORF3 and ORF4 specify a glutathione S-transferase and a polypeptide showing no homology to known proteins, respectively.

rified and shown by ultraviolet-visible absorption spectrophotometry and EPR spectroscopy to contain a putidaredoxin-type [2Fe-2S] cluster (4). The redox potential of the cluster was determined to be $-245 (\pm 5)$ mV, some 100 mV lower than the values observed for electron shuttles acting with Class IIB or Class III dioxygenases (44, 62).

A similar strategy was employed to amplify part of the gene of the reductase component of the electron supply system. In this case, the degenerate primers were designed from the N-terminal sequence of the protein (13) and from a well-conserved motif assumed to be located at its C-terminus. The probe obtained was used to screen the cosmid library of RW1 DNA and a large DNA fragment encompassing the reductase gene was identified and characterized. Comparison of the predicted amino acid sequence of the reductase with sequences in the EMBL/GENBANK/DDBJ data banks revealed some marked similarities with putidaredoxin reductase (42, 54), terpredoxin reductase (53) and rhodocoxin reductase (48), and some less well defined similarities with reductases acting with Class IIB dioxygenases. Few similarities with reductases of other classes of multicomponent non-heme iron mono- and dioxygenases were found, which is consistent with the biochemical and functional characteristics of these proteins: the stereospecificity of removal of the hydride ion from NADH by Class II A and B reductases are *pro*-S whereas all others remove the *pro*-R hydride ion (57). The cloned reductase gene was hyperexpressed in *E. coli* and the resulting reductase was purified and shown to donate electrons to the *fdx1* ferredoxin in a cytochrome c reduction assay. The availability of the two genes specifying the electron supply system of the dioxin dioxygenase will permit in the near future the construction of optimized cassettes for the heterologous expression of functional dioxin dioxygenase.

The genes specifying the different components of the initial dioxygenase are not clustered

The genes specifying the different polypeptides of three-component dioxygenase systems (Class IIB and Class III) are usually clustered (Fig. 4). In some cases, such as in the polychlorinated biphenyl degrader *Pseudomonas* sp. KKS102, the reductase gene is not contiguous with the three other genes, but nevertheless present in the same locus (39). The localization of the ferredoxin and the reductase genes on different DNA fragments, neither of which encodes the dioxin dioxygenase (4), is thus unusual. Whether or not this unusual genetic organization will turn out to be common for Class IIA ring-hydroxylating dioxygenases, whether *Sphingomonas* sp. RW1 generally does not have its catabolic determinants clustered in operons, or whether such genetic organization is rare and limited to the dioxin catabolic pathway, remain to be determined.

Earlier unsuccessful attempts to clone the dioxin dioxygenase genes by constructing an expression library and screening for ring-hydroxylating activity are now explained as there is no linkage of the genes encoding the components of the dioxygenase. Moreover, problems to coexpress a similar ferredoxin and its specific reductase have been experienced (54). The presence of both putidaredoxin reductase and putidaredoxin in the same *E. coli* cell line might perhaps result in cell death due to leakage of electrons from NADH through the reductase to putidaredoxin, which has a greater sensitivity to oxidation by molecular oxygen than do the other components of this electron transfer system. Therefore, efforts to co-express all components of the dioxygenase may need to rely on the use of an inducible expression system.

The gene encoding the *meta*-cleavage enzyme is also not directly linked to the dioxin dioxygenase genes

The second step of the dioxin catabolic pathway, the cleavage of the dihydroxylated aromatic ring by the introduction of two further oxygen atoms, is carried out by DbfB, a *meta*-cleavage enzyme, 2,2',3-trihydroxybiphenyl-1,2-dioxygenase (Fig. 1). The structural gene coding for this enzyme has been cloned from a cosmid expression library using a functional colorimetric assay, and sequenced (34). The *dbfB* gene does not appear to be directly linked with the dioxin dioxygenase genes nor with the genes specifying the other enzymes of the path-

ways. The amino acid sequence of this enzyme is typical of those of extradiol dioxygenases, as was shown recently by a structure-validated alignment (20, 33). After hyperexpression of its structural gene in *E. coli*, the protein has been purified anaerobically and characterized. 2,2',3-trihydroxybiphenyl and 2,3-dihydroxybiphenyl are substrates of the enzyme and, like the substrates of many *meta*-cleavage enzymes (19), inhibit the enzyme when present at high concentration (34). Although the size of the polypeptide is close to that of other *meta*-cleavage enzymes, it was found to be active as a monomer in contrast to all other extradiol dioxygenases thus far studied which are active as homotetrameric or homooctameric proteins (34). This feature of DbfB makes nuclear magnetic resonance studies feasible and has allowed identification of the ligands of ferrous iron in the catalytic centre of this enzyme (10).

RW1 has been shown to contain at least three *meta*-cleavage enzymes (13, 34) and another dibenzofuran degrader, *Terrabacter* sp. strain DPO 360, also contains several distinct extradiol dioxygenases (58). One PCB-degrader, *Rhodococcus globerulus* P6, has been shown to possess up to eight different *meta*-cleavage enzymes (6, 43). The reason for the multiplicity of such isoenzymes has not yet been definitively explained. While the protein DbfB is thought to be the *meta*-cleavage enzyme involved in the "upper" pathway of dibenzo-*p*-dioxin and dibenzofuran degradation, the two other proteins may be involved in the ring-cleavage of catechol or chlorocatechol of the *meta* pathway. One of the isoenzymes of P6 has been shown to be less sensitive to substrate-mediated inactivation by chlorinated substrates, suggesting that it may be important for the catabolism of certain chlorinated congeners (Prucha, M. and K.N. Timmis, unpublished results).

The hydrolytic step of the dibenzofuran pathway

Although the ring cleavage product in the dioxin pathway, 2-hydroxy-6-oxo-6-(2-hydroxyphenoxy)-hexa-2,4-dienoate, spontaneously decomposes to catechol and hydroxy-*cis,cis*-muconate, the corresponding product in the dibenzofuran pathway, 2-hydroxy-6-oxo-6-(2-hydroxyphenyl)-hexa-2,4-dienoate (2-OH-HOPDA), is enzymatically hydrolysed to salicylate and 2-hydroxypentadienoate (Fig. 1). Bünz *et al.* (14) have purified from *Sphingomonas* sp. RW1 two different hydrolases able to convert the substrate 2-OH-HOPDA. The N-terminal sequences of these two hydrolases are quite

different from known hydrolases. However, the two proteins are isoenzymes and exhibit 50% similarity within their 20 first amino acids. Such diversity of hydrolases which apparently have the same substrate range, if not the same specific activity (14), is unusual. The reason is unclear but possibly one of the hydrolases play a role in a second degradative pathway, such as for biphenyl since biphenyl-utilizing mutants of this organism arise spontaneously (71). Preliminary data indicate that the RW1 strain may contain at least four different genes encoding such hydrolases (D'Enza, M., B. Happe, and J. Armengaud, unpublished results). It is known that different hydrolases exhibit different specificities towards bicyclic or monocyclic aromatic substrates (28) and, it will thus be of interest to study these isoenzymes in terms of substrate range and kinetic parameters.

Conclusions

The present review highlights the atypical genetic organization of the degradative pathways of dibenzo-*p*-dioxin, dibenzofuran and their chloro-derivatives in *Sphingomonas* sp. RW1, which seem to be significantly different from those of other aromatic compounds, such as naphthalene, cumene, toluene, benzene and biphenyl. The recent cloning of the genes specifying the enzymatic components of the enzymes of the initial steps in the dibenzofuran and dibenzo-*p*-dioxin pathways, and analysis of the genes and their products, have opened new perspectives for the degradation of such compounds and represent an important base from which to obtain a more complete description of the genetic, biochemical and regulational levels of the different actors involved in the catabolism of these compounds. This will provide the knowledge base for development of rational strategies for improving catabolic activities towards chlorinated-derivatives, and for bioremediation applications.

Acknowledgments

We gratefully acknowledge our numerous colleagues for their important contributions to the RW1 pathways, and in particular B. Averhoff, M. D'Enza, B. Happe, G. Manicini, D. Pieper, H. Poth, M. Strätz, R.-M. Wittich, and our collaboration partners at the University of Hamburg.

J. Armengaud was supported by a long-term grant from the FEBS and wishes to repeat his deep gratitude to the persons acting generously in this federation. This research was supported by the German Ministry for Education and Research

(BMBF grant 0318896C). K. N. Timmis gratefully acknowledges the generous support of the Fonds der Chemischen Industrie.

References

1. **Abramowicz, D.A.** 1990. Aerobic and anaerobic biodegradation of PCBs: a review. *Crit. Rev. Biotechnol.* **10**, 241-251.
2. **Arfmann, H.-A., K.N. Timmis, and R.-M. Wittich.** 1997. Mineralization of 4-chlorodibenzofuran by a consortium consisting of *Sphingomonas* sp. strain RW1 and *Burkholderia* sp. strain JWS. *Appl. Environ. Microbiol.* **63**, 3458-3462.
3. **Armengaud, J., C. Meyer, and Y. Jouanneau.** 1997. A [2Fe-2S] ferredoxin (FdVI) is essential for growth of the photosynthetic bacterium *Rhodobacter capsulatus*. *J. Bacteriol.* **179**, 3304-3309.
4. **Armengaud, J. and K.N. Timmis.** 1997. Molecular characterization of *Fdx1*, a putidaredoxin-type [2Fe-2S] ferredoxin able to transfer electrons to the dioxin dioxygenase of *Sphingomonas* sp. RW1. *Eur. J. Biochem.* **247**, 833-842.
5. **Asturias, J.A., E. Diaz, and K.N. Timmis.** 1995. The evolutionary relationship of biphenyl dioxygenase from Gram-positive *Rhodococcus globerulus* P6 to multicomponent dioxygenases from Gram-negative bacteria. *Gene* **156**, 11-18.
6. **Asturias, J.A. and K.N. Timmis.** 1993. Three different 2,3-dihydroxybiphenyl-1,2-dioxygenase genes in the gram-positive polychlorobiphenyl-degrading bacterium *Rhodococcus globerulus* P6. *J. Bacteriol.* **175**, 4631-4640.
7. **Asturias, J.A., L.D. Eltis, M. Prucha, and K.N. Timmis.** 1994. Analysis of three 2,3-dihydroxybiphenyl 1,2-dioxygenases found in *Rhodococcus globerulus* P6. Identification of a new family of extradiol dioxygenases. *J. Biol. Chem.* **269**, 7807-7815.
8. **Ayres, D.C.** 1981. Destruction of polychlorodibenzo-*p*-dioxins. *Nature* **290**, 323-324.
9. **Batie, C.J., D.P. Ballou, and C.C. Correll.** 1992. Phthalate dioxygenase reductase and related flavin-iron-sulfur containing electron transferases. p543-566. In F. Müller (ed.). Chemistry and Biochemistry of flavoenzymes. vol 3. CRC Press. Boca Raton, Fla.
10. **Bertini, I., F. Capozzi, A. Dikiy, B. Happe, C. Luchinat, and K.N. Timmis.** 1995. Evidence of histidine coordination to the catalytic ferrous ion in the ring-cleaving 2,2',3-trihydroxybiphenyl dioxygenase from the dibenzofuran-degrading bacterium *Sphingomonas* sp. strain RW1. *Biochem. Biophys. Res. Commun.* **215**, 855-860.
11. **Beurskens, J.E.M., M. Toussaint, J. De Wolf, J.M.D. Van der Steen, P.C. Slot, L.C.M. Commandeur, and J.R. Parsons.** 1995. Dehalogenation of chlorinated dioxins by an anaerobic microbial consortium from sediment. *Environ. Toxicol. Chem.* **14**, 939-943.
12. **Bruschi, M. and F. Guerlesquin.** 1988. Structure, function and evolution of bacterial ferredoxins. *FEMS Microbiol. Rev.* **54**, 155-176.
13. **Bünz, P.V. and A.M. Cook.** 1993. Dibenzofuran 4,4a-dioxygenase from *Sphingomonas* sp. strain RW1: angular dioxygenation by a three-component enzyme system. *J. Bacteriol.* **175**, 6467-6475.
14. **Bünz, P.V., R. Falchetto, and A.M. Cook.** 1993. Purification of two isofunctional hydrolases (EC 3.7.1.8) in the degradative pathway for dibenzofuran in *Sphingomonas* sp. strain RW1. *Biodegradation* **4**, 1171-1178.
15. **Butler, C.S. and J.R. Mason.** 1997. Structure-function analysis of the bacterial aromatic ring-hydroxylating dioxygenases. *Adv. Microb. Physiol.* **38**, 47-84.
16. **Chang, C.Y. D.A. Wu, C.C. Lai, W.L. Miller, and B.C. Chung.** 1988. Cloning and structure of the human adrenodoxin gene. *DNA* **7**, 609-15.
17. **Denome, S.A., D.C. Stanley, E.S. Olson, and K.D. Young.** 1993. Metabolism of dibenzothiophene and naphthalene in *Pseudomonas* strains: complete DNA sequence of an upper naphthalene catabolic pathway. *J. Bacteriol.* **175**, 6890-6901.
18. **Dobbs, A.J. and C. Grant.** 1979. Photolysis of highly chlorinated dibenzo-*p*-dioxins by sunlight. *Nature* **278**, 163-165.
19. **Eltis, L.D., B., Hofmann, H.-J. Hecht, H. Lnsdorf, and K.N. Timmis.** 1993. Purification and crystallisation of 2,3-dihydroxybiphenyl 1,2-dioxygenase. *J. Biol. Chem.* **268**, 2727-2732.
20. **Eltis, L.D. and J.T. Bolin.** 1996. Evolutionary relationships among extradiol dioxygenases. *J. Bacteriol.* **178**, 5930-5937.
21. **Engesser, K.H., V. Strubel, K. Christoglou, P. Fischer, and H.G. Rast.** 1989. Dioxygenolytic cleavage of aryl ether bonds: 1,10-dihydro-1,10-dihydroxyfluoren-9-one, a novel arene dihydrodiol as evidence for angular dioxygenation of dibenzofuran. *FEMS Microbiol. Lett.* **53**, 205-209.
22. **Ensley, B.D., D.T. Gibson, and A.L. LaBorde.** 1982. Oxidation of naphthalene by a multicomponent enzyme system from *Pseudomonas* sp. strain NCIB 9816. *J. Bacteriol.* **149**, 948-954.
23. **Erickson, B.D. and F.J. Mondello.** 1992. Nucleotide sequencing and transcriptional mapping of the genes encoding biphenyl dioxygenase, a multicomponent polychlorinated-biphenyl-degrading enzyme in *Pseudomonas* strain LB400. *J. Bacteriol.* **174**, 2903-2912.
24. **Erickson, B.D. and F.J. Mondello.** 1993. Enhanced biodegradation of polychlorinated biphenyls after site-directed mutagenesis of a biphenyl dioxygenase gene. *Appl. Environ. Microbiol.* **59**, 3858-3862.
25. **Fletcher C.L. and W.A. McKay.** 1993. Polychlorinated dibenzo-*p*-dioxins (PCDDs) and dibenzofurans (PCDFs) in the aquatic environment-a literature review. *Chemosphere* **26**, 1041-1069.
26. **Fortnagel, P., H. Harms, R.-M. Wittich, S. Krohn, H. Meyer, V. Sinnwell, H. Wilkes, and W. Francke.** 1990. Metabolism of dibenzofuran by *Pseudomonas* sp. strain HH69 and the mixed culture HH27. *Appl. Environ. Microbiol.* **56**, 1148-1156.
27. **Furukawa, K.** 1994. Molecular genetics and evolutionary relationship of PCB-degrading bacteria. *Biodegradation* **5**, 289-300.
28. **Furukawa, K., J. Hirose, A. Suyama, T. Zaiki, and S. Hayashida.** 1993. Gene components responsible for discrete substrate specificity in the metabolism of biphenyl (*bph* operon) and toluene (*tod* operon). *J. Bac-*

- teriol. **175**, 5224-5232.
29. **Furukawa, K., N. Arimura, and T. Miyazaki.** 1987. Nucleotide sequence of the 2,3-dihydroxybiphenyl dioxygenase gene of *Pseudomonas pseudoalcaligenes*. *J. Bacteriol.* **169**, 427-429.
 30. **Gibson, D.T. and V. Subramanian.** 1984. Microbial degradation of aromatic hydrocarbons. p. 181-252. In D.T. Gibson (ed.). Microbial degradation of organic compounds, Marcel Dekker, New York, NY.
 31. **Gray, K. O.S. Pogrebinsky, G.T. Mrachko, L. Xi, D.J. Montecillo, and C.H. Squires.** 1996. Molecular mechanisms of biocatalytic desulfurization of fossil fuels. *Nature Biotechnology* **14**, 1705-1709.
 32. **Haddock, J.D., J.R. Horton, and D.T. Gibson.** 1995. Dihydroxylation and dechlorination of chlorinated biphenyls by purified biphenyl 2,3-dioxygenase from *Pseudomonas* sp. strain LB400. *J. Bacteriol.* **177**, 20-26.
 33. **Han, S., L.D. Eltis, K.N. Timmis, S.W. Muchmore, and J.T. Bolin.** 1995. Crystal structure of the biphenyl-cleaving extradiol dioxygenase from a PCB-degrading *Pseudomonas*. *Science* **270**, 976-980.
 34. **Happe, B., L.D. Eltis, H. Poth, R. Hedderich, and K. N. Timmis.** 1993. Characterization of 2,2',3-trihydroxybiphenyl dioxygenase, an extradiol dioxygenase from the dibenzofuran- and dibenzo-*p*-dioxin-degrading bacterium *Sphingomonas* sp. strain RW1. *J. Bacteriol.* **175**, 7313-7320.
 35. **Harms, H., R.-M. Wittich, V. Sinnwell, H. Meyer, P. Fortnagel, and W. Francke.** 1990. Transformation of dibenzo-*p*-dioxin by *Pseudomonas* sp. strain HH69. *Appl. Environ. Microbiol.* **56**, 1157-1159.
 36. **Hay, A.** 1979. Séveso: the crucial question of reactor safety. *Nature* **281**, 521.
 37. **Hofer, B., D. Eltis, D.N. Dowling, and K.N. Timmis.** 1993. Genetic analysis of a *Pseudomonas* locus encoding a pathway for biphenyl/polychlorinated biphenyl degradation. *Gene* **130**, 47-55.
 38. **Hofer, B., S. Backhaus, and K.N. Timmis.** 1994. The biphenyl/polychlorinated biphenyl-degradation locus (*bph*) of *Pseudomonas* sp. LB400 encodes four additional metabolic enzymes. *Gene* **144**, 9-16.
 39. **Kikuchi, Y., Y. Nagata, M. Hinata, K. Kimbara, M. Fukuda, K. Yano, and M. Takagi.** 1994. Identification of the *bphA4* gene encoding ferredoxin reductase involved in biphenyl and polychlorinated biphenyl degradation in *Pseudomonas* sp. strain KKS102. *J. Bacteriol.* **176**, 1689-1694.
 40. **Kimbara, K., T. Hashimoto, M. Fukuda, T. Koana, M. Takagi, M. Oishi, and K. Yano.** 1989. Cloning and sequencing of two tandem genes involved in degradation of 2,3-dihydroxybiphenyl to benzoic acid in the polychlorinated biphenyl-degrading soil bacterium *Pseudomonas* sp. strain KKS102. *J. Bacteriol.* **171**, 2740-2747.
 41. **Kimura, N., A. Nishi, M. Goto, and K. Furukawa.** 1997. Functional analyses of a variety of chimeric dioxygenases constructed from two biphenyl dioxygenases that are similar structurally but different functionally. *J. Bacteriol.* **179**, 3936-3943.
 42. **Koga, H., E. Yamaguchi, K. Matsunaga, H. Aramaki, and T. Horiuchi.** 1989. Cloning and nucleotide sequences of NADH-putidaredoxin reductase gene (*camA*) and putidaredoxin gene (*camB*) involved in cytochrome P-450cam hydroxylase of *Pseudomonas putida*. *J. Biochem.* **106**, 831-836.
 43. **Kudo, T., M. Maeda, and F. Fuji.** 1996. Diversity of *bph* gene cluster of PCB-degrading Rhodococci. p47. In K.N. Timmis and J. Lalucat (ed.) "Biodegradation of organic pollutants" abstract book, UIB-GBF-CSIC-TUB symposium, Mallorca 1996.
 44. **Link, T.A., O.M. Hatzfeld, P. Unalkat, J.K. Shergill, R. Cammack, and J.R., Mason.** 1996. Comparison of the "Rieske" [2Fe-2S] center in the bc1 complex and in bacterial dioxygenases by circular dichroism spectroscopy and cyclic voltammetry. *Biochemistry* **35**, 7546-7552.
 45. **Megharaj, M., R.-M. Wittich, R. Blasco, D.H. Pieper, and K.N. Timmis.** 1997. Superior survival and degradation of dibenzo-*p*-dioxin and dibenzofuran in soil by soil-adapted *Sphingomonas* sp. strain RW1. *Appl. Microbiol. Biotechnol.* **48**, 109-114.
 46. **Mohn, W.W. and J.M., Tiedje.** 1992. Microbial reductive dehalogenation. *Microbiol. Rev.* **56**, 285-289.
 47. **Monna, L., T. Omori, and T. Kodama.** 1993. Microbial degradation of dibenzofuran, fluorene and dibenzo-*p*-dioxin by *Staphylococcus auricularis* DBF63. *Appl. Environ. Microbiol.* **59**, 285-289.
 48. **Nagy, I., G. Schoofs, F. Compennolle, P. Proost, J. Vanderleyden, and R. de Mot.** 1995. Degradation of the thiocarbamate herbicide EPTC (S-ethyl dipropylcarbamothioate) and biosafening by *Rhodococcus* sp. strain NI86/21 involve an inducible cytochrome P-450 system and aldehyde dehydrogenase. *J. Bacteriol.* **177**, 676-687.
 49. **Nakatsu, C.H., N.A. Straus, and R.C. Wyndham.** 1995. The nucleotide sequence of the Tn5271 3-chlorobenzoate 3, 4-dioxygenase genes (*cbaAB*) unites the class IA oxygenases in a single lineage. *Microbiology* **141**, 485-495.
 50. **Naud, I., M. Vinon, J. Garin, J. Gaillard, E., Forest, and Y. Jouanneau.** 1994. Purification of a sixth ferredoxin from *Rhodobacter capsulatus*: primary structure and biochemical properties. *Eur. J. Biochem.*, **222**, 933-939.
 51. **Neidle, E.L., C. Hartnett, L.N. Ornston, A. Bairoch, M. Rekik, and S. Harayama.** 1992. *Cis*-diol dehydrogenases encoded by the TOL pWW0 plasmid *xylL* gene and the *Acinetobacter calcoaceticus* chromosomal *benD* gene are members of the short-chain alcohol dehydrogenase superfamily. *Eur. J. Biochem.* **204**, 113-120.
 52. **Parales, J.V., A. Kumar, R.E. Parales, and D.T. Gibson.** 1996. Cloning and sequencing of the genes encoding 2-nitrotoluene dioxygenase from *Pseudomonas* sp. JS42. *Gene* **181**, 57-61.
 53. **Peterson, J.A., J.Y. Lu, J. Geisselsoder, S. Graham-Lorence, C. Carmona, F. Witney, and M.C. Lorence.** 1992. Cytochrome P-450terp. Isolation and purification of the protein and cloning and sequencing of its operon. *J. Biol. Chem.* **267**, 14193-14203.
 54. **Peterson, J.A., M.C. Lorence, and B. Amarneh.** 1990. Putidaredoxin reductase and putidaredoxin. Cloning, sequence determination and heterologous expression of the proteins. *J. Biol. Chem.* **265**, 6066-6073.
 55. **Sariaslani, F.S.** 1989. Microbial enzymes for oxidation

- of organic molecules. *CRC Crit. Rev. Biotech.* **9**, 171-257.
56. **Sauber, K., C. Fröhner, G. Rosenberg, J. Eberspacher, and F. Lingens.** 1977. Purification and properties of pyrazon dioxygenase from pyrazon-degrading bacteria. *Eur. J. Biochem.* **74**, 89-97.
 57. **Schläfli, H.R., D.P. Baker, T. Leisinger, and A.M. Cook.** 1995. Stereospecificity of hydride removal from NADH by reductases of multicomponent nonheme iron oxygenase systems. *J. Bacteriol.* **177**, 831-834.
 58. **Schmid, A., B. Rothe, J. Altenbuchner, W. Ludwig, and K.-H. Engesser.** 1997. Characterization of three distinct extradiol dioxygenases involved in mineralization of dibenzofuran by *Terrabacter* sp. strain DPO360. *J. Bacteriol.* **179**, 53-62.
 59. **Schmid, A., H.-P. E. Kohler, J.-R. Van der Meer, H.-J. Knackmuss, and K.-H. Engesser.** 1996. Bacterial mineralization of 2-hydroxybiphenyl: a link between phenol-, dibenzofuran- and biphenyl-degradation pathways. p.163. In K.N. Timmis and J. Lalucat (ed.) "Biodegradation of organic pollutants" abstract book, UIB-GBF-CSIC-TUB symposium, Mallorca 1996.
 60. **Strubel, V., H. G. Rast, W. Fietz, H.-J. Knackmuss, and K.H. Engesser.** 1989. Enrichment of dibenzofuran utilizing bacteria with high co-metabolic potential towards dibenzodioxin and other anellated aromatics. *FEMS Microbiol. Lett.* **58**, 233-238.
 61. **Strubel, V., K.-H. Engesser, P. Fischer, and H.-J., Knackmuss.** 1991. 3-(2-Hydroxyphenyl)catechol as substrate for proximal meta ring cleavage in dibenzofuran degradation by *Brevibacterium* sp. strain DPO 1361. *J. Bacteriol.* **173**, 1932-1937.
 62. **Subramanian, V., T.-N. Liu, W.-K. Yeh, C.M. Serdar, L.P. Wackett, and D.T., Gibson.** 1985. Purification and properties of FerredoxinTOL, a component of toluene dioxygenase from *Pseudomonas putida* F1. *J. Biol. Chem.* **260**, 2355-2363.
 63. **Suyama, A., R. Iwakiri, N. Kimura, A. Nishi, K. Nakamura, and K. Furukawa.** 1996. Engineering hybrid *Pseudomonads* capable of utilizing a wide range of aromatic hydrocarbons and of efficient degradation of trichloroethylene. *J. Bacteriol.* **178**, 4039-4046.
 64. **Sylvestre, M., M. Sirois, Y. Hurtubise, J. Bergeron, D. Ahmad, F. Shareck, D. Barriault, I. Guillemette, and J.M. Juteau.** 1996. Sequencing of *Comamonas testosteroni* strain B-356-biphenyl/chlorobiphenyl dioxygenase genes: evolutionary relationships among Gram-negative bacterial biphenyl dioxygenases. *Gene* **174**, 195-202.
 65. **Ta, D.T. and L.E. Vickery.** 1992. Cloning, sequencing and overexpression of a [2Fe-2S] ferredoxin gene from *Escherichia coli*. *J. Biol. Chem.* **267**, 11120-11125.
 66. **Tiedje, J.M., J.F. 3rd. Quensen, J. Chee-Sanford, J. P. Schimel, and S.A. Boyd.** 1993. Microbial reductive dechlorination of PCBs. *Biodegradation* **4**, 231-240.
 67. **Van den Berg, M., J. De Jongh, H. Poiger, and J.R. Olson.** 1994. The toxicokinetics and metabolism of polychlorinated dibenzo-*p*-dioxins (PCDDs) and dibenzofurans (PCDFs) and their relevance for toxicity. *Crit. Rev. Toxicol.* **24**, 1-74.
 68. **Vuilleumier, S.** 1997. Bacterial glutathione S-transferases: what are they good for? *J. Bacteriol.* **179**, 1431-1441.
 69. **Wang, S.P., Y.P. Chen, and B. Ely.** 1995. A ferredoxin, designed FdxP, stimulates *p*-hydroxybenzoate hydroxylase activity in *Caulobacter crescentus*. *J. Bacteriol.* **177**, 2908-2911.
 70. **Wilkes, H., R.-M., Wittich, K.N. Timmis, P. Fortnagel, and W. Francke.** 1996. Degradation of chlorinated dibenzofurans and dibenzo-*p*-dioxins by *Sphingomonas* sp. strain RW1. *Appl. Environ. Microbiol.* **62**, 367-371.
 71. **Wittich, R.-M., H. Wilkes, V. Sinnwell, W. Francke, and P. Fortnagel.** 1992. Metabolism of dibenzo-*p*-dioxin by *Sphingomonas* sp. strain RW1. *Appl. Environ. Microbiol.* **58**, 1005-1010.
 72. **Zylstra, G.J. and D.T. Gibson.** 1989. Toluene degradation by *Pseudomonas putida* F1. Nucleotide sequence of the *TodC1C2BADE* genes and their expression in *Escherichia coli*. *J. Biol. Chem.* **264**, 14940-14946.
 73. **Zylstra, G.J. and D.T. Gibson.** 1991. Aromatic hydrocarbon degradation: a molecular approach. *Genetic Engineering* **13**, 183-203.